

Stability analysis of the Randall-Sundrum braneworld in presence of bulk scalar

Debaprasad Maity*, Soumitra SenGupta †

*Department of Theoretical Physics,
Indian Association for the Cultivation of Science,
Calcutta - 700 032, India*

Sourav Sur‡

Institute of Physics, Bhubaneswar - 751 005, India

The stability problem of Randall-Sundrum braneworld is readdressed in the light of stabilizing bulk scalar fields. It is shown that in such scenario the instability persists because of back-reaction even when an arbitrary potential is introduced for a canonical scalar field in the bulk. It is further shown that a bulk scalar field can indeed stabilize the braneworld when it has a tachyon-like action. The full back-reacted metric in such model is derived and a proper resolution of the hierarchy problem (for which the Randall Sundrum scenario was originally proposed) is found to exist by suitable adjustments of the parameters of the scalar potential.

Standard model for strong, weak and electromagnetic interactions based on the gauge group $SU(3) \times SU(2) \times U(1)$ has been extremely successful in explaining physical phenomena upto TeV scale. Such a model, however, encounters the well known fine tuning problem in connection with Higgs mass related to the gauge hierarchy problem which refers to the vast disparity between the weak and Planck scale. By invoking supersymmetry one can resolve this problem at the expense of incorporating a large number of (hitherto unseen) superpartners in the theory. In an alternative approach, theories with extra spatial dimension(s) have attracted a lot of attention because of the new geometric approach to solve the same problem and also the distinct features from ordinary Kaluza-Klein theory. In such models the standard model fields are localized on a 3+1 dimensional brane [1, 2, 3, 4, 5, 6, 7] where gravity can propagate in the bulk spacetime. One of the most theoretically appealing and phenomenologically interesting model in this context was proposed by Randall and Sundrum (RS) [4] where the mass hierarchy emerged naturally in an exponentially warped geometry along the extra dimension. The model contains two 3+1 dimensional branes sitting at the two orbifold fixed points where the single extra dimension in a 4+1 dimensional bulk has been compactified on a S_1/Z_2 orbifold. The model contains two parameters namely the bulk cosmological constant Λ and the brane separation r_c . The geometry and the Higgs mass is warped exponentially by a dimensionless parameter kr_c , where $k = \sqrt{-\Lambda/24M^3}$, M being the 5D Planck mass. For the desired warping one should have $kr_c \sim 11$. If $k \sim$ Planck mass then, r_c should have a stable value near Planck length. To stabilize the value of r_c , Goldberger and Wise (GW) [8] proposed a simple mechanism by introducing a minimally coupled massive scalar field in the bulk. Later several other works have been done in this

direction [9, 10, 11, 12]. In the original work of GW the effect of back-reaction of the scalar field on the background metric was neglected. Such back-reaction was later included in subsequent works and a modified solution for the metric was found [13, 14]. In this paper we carefully re-examine the stability issue in the back-reacted RS model when the scalar field is introduced in bulk. We show on a very general ground that stabilization is not possible with a minimally coupled scalar field in the bulk. Following GW we subsequently show that if the dependence of scalar field action on the extra coordinate is tachyon-like then we can stabilize the brane world and the stabilized value of r_c can produce the desired hierarchy from Planck scale to TeV scale.

Let us first consider the following bulk action [8, 13, 14, 15]

$$S = \int d^5x \sqrt{-G} \left[-M^3 R + \sum_{i=1}^n \left(\frac{1}{2} \partial_A \phi_i \partial^A \phi_i \right) - V(\{\phi_i\}) \right] - \int d^4x dy \sqrt{-g_a} \delta(y - y_a) \lambda_a(\{\phi_i\}) \quad (1)$$

where $\{\phi_i\} = \{\phi_1 \cdots \phi_n\}$ are a set of n scalar fields with a mixed potential $V(\{\phi_i\})$ and the index a runs over the brane locations. The corresponding brane potentials are denoted by λ_a .

Taking the line element in the form

$$ds^2 = e^{-2A(y)} \eta_{\mu\nu} dx^\mu dx^\nu - dy^2 \quad (2)$$

the field equations are given as

$$\phi_i'' - 4A'\phi_i' = \frac{\partial V}{\partial \phi_i} + \sum_a \frac{\partial \lambda_a}{\partial \phi_i} \delta(y - y_a) \quad (3a)$$

$$4A'^2 - A'' = -\frac{V}{3M^3} - \frac{1}{6M^3} \sum_a \lambda_a \delta(y - y_a) \quad (3b)$$

$$4A'^2 - 4A'' = -\frac{V}{3M^3} - \frac{1}{2M^3} \sum_i \phi_i'^2 - \frac{2}{3M^3} \sum_a \lambda_a \delta(y - y_a) \quad (3c)$$

*E-mail: tpdm@mahendra@iacs.res.in

†E-mail: tpssg@mahendra@iacs.res.in

‡E-mail: sourav@iopb.res.in

where prime $\{\prime\}$ denotes partial differentiation with respect to the extra spatial coordinate y .

The boundary conditions are

$$[\phi'_i]_a = \frac{\partial \lambda_a}{\partial \phi_i}; \quad [A']_a = \frac{1}{6M^3} \lambda_a(\{\phi_i\}). \quad (4)$$

Denoting $C = 1/24M^3$, it follows

$$A'^2 = 2C \left(\sum_i \phi_i'^2 - 2V \right); \quad A'' = 4C \sum_i \phi_i'^2 \quad (5a)$$

$$\phi_i'' = 4A' \phi_i' + \frac{\partial V}{\partial \phi_i} \quad (5b)$$

In order to obtain analytic closed form solutions we resort to the particular class of effective potentials as in refs. [13, 16]:

$$V(\{\phi_i\}) = \frac{1}{8} \sum_{i=1}^n \left[\frac{\partial W}{\partial \phi_i} \right]^2 - 2CW^2 \quad (6)$$

where

$$W(\{\phi_i\}) = \sum_{i=1}^n W_i(\phi_i) \quad (7)$$

is a sum of n superpotentials $\{W_1(\phi_1), \dots, W_n(\phi_n)\}$.

Eqs.(5) lead to the first order equations:

$$\phi_i' = \frac{1}{2} \frac{\partial W}{\partial \phi_i}; \quad A' = 2CW. \quad (8)$$

Following GW [8], we calculate the effective potential V_{eff} on the 3-brane as:

$$V_{eff} = 2 \int_0^{y_\pi} dy e^{-4A(y)} \left[\frac{1}{2} \sum_{i=1}^n \partial_A \phi_i \partial^A \phi_i - V(\{\phi_i\}) \right] - e^{-4A(0)} \lambda_0(\{\phi_i^0\}) - e^{-4A(y_\pi)} \lambda_\pi(\{\phi_i^\pi\}) \quad (9)$$

where the Planck brane is located at $y = 0$ and the visible brane is at $y = y_\pi$, and we have used the notations: $\phi_i^0 \equiv \phi_i(y = 0)$, $\phi_i^\pi \equiv \phi_i(y = y_\pi)$ for $i = 1, \dots, n$. Using the boundary conditions (4) which now take the form

$$[W(\{\phi_i\})]_0 = 2\lambda_0(\{\phi_i\}); \quad \left[\frac{\partial W}{\partial \phi_i} \right]_0 = 2 \frac{\partial \lambda_0}{\partial \phi_i} \quad (10a)$$

$$[W(\{\phi_i\})]_{y_\pi} = 2\lambda_\pi(\{\phi_i\}); \quad \left[\frac{\partial W}{\partial \phi_i} \right]_{y_\pi} = 2 \frac{\partial \lambda_\pi}{\partial \phi_i} \quad (10b)$$

the brane potentials $\lambda_0(\phi)$ and $\lambda_\pi(\phi)$ are given by [13]

$$\lambda_0(\{\phi_i\}) = W(\{\phi_i^0\}) + \sum_{i=1}^n \left[\frac{\partial W}{\partial \phi_i^0} (\phi_i - \phi_i^0) \right] + \sum_{i=1}^n \gamma_{0i}^2 (\phi_i - \phi_i^0)^2 \quad (11a)$$

$$\lambda_\pi(\{\phi_i\}) = -W(\{\phi_i^\pi\}) - \sum_{i=1}^n \left[\frac{\partial W}{\partial \phi_i^\pi} (\phi_i - \phi_i^\pi) \right] + \sum_{i=1}^n \gamma_{\pi i}^2 (\phi_i - \phi_i^\pi)^2 \quad (11b)$$

where the constants γ_{0i} and $\gamma_{\pi i}$, (for $i = 1, \dots, n$) are parameters of various potentials.

Now, in order to have extremum for the above effective potential at some value of y_π ($= y_s$, say)

$$\left[\frac{\partial V_{eff}}{\partial y_\pi} \right]_{y_\pi=y_s} = -4C e^{-4A(y_s)} W(y_s)^2 = 0. \quad (12)$$

which implies $W(y_\pi) = 0$ as well as $\partial^2 V_{eff} / \partial y_\pi^2 = 0$ at $y_\pi = y_s$. On the other hand, the third derivative is given by

$$\left[\frac{\partial^3 V_{eff}}{\partial y_\pi^3} \right]_{y_\pi=y_s} = -4C e^{-4A(y_s)} \left[\frac{\partial W(y_\pi)}{\partial y_\pi} \right]_{y_\pi=y_s}^2 \quad (13)$$

which is clearly non-vanishing in accord with the above boundary conditions. Therefore, no extremum exists and we only have a point of inflection at the value of $y_\pi = y_s$ for which W vanishes.

Let us now resort to a general solution ansatz for the metric function $A(y)$ and single scalar field ϕ so that the standard RS [4] solution is obtained in the limit where the scalar $\phi(y)$ becomes trivial:

$$A(y) = ky + f(\phi_{cl}(y)); \quad \phi = \phi_{cl}(y, a_1, a_2) \quad (14)$$

where a_1, a_2 are the initial parameters of the theory and constant $k \sim 5D$ Planck scale as in the RS picture.

The field equations (5) yield

$$\phi_{cl}'^2 = \frac{1}{4C} \frac{\partial^2 f}{\partial y^2} \quad (15a)$$

$$V = \frac{1}{8C} \frac{\partial^2 f}{\partial y^2} - \frac{1}{2C} \left(k + \frac{\partial f}{\partial y} \right)^2. \quad (15b)$$

In the limit $f \rightarrow 0$, $V \rightarrow -12M^3 k^2 = \Lambda$ which gives the standard AdS_5 bulk geometry of RS model.

From Eqs.(4), the boundary values of the brane potentials and their first derivatives are obtained as

$$[\lambda_0]_0 = 2C \left[k + \frac{\partial f}{\partial y} \right]_0; \quad \left[\frac{\partial \lambda_0}{\partial y} \right]_0 = 2C \left[\frac{\partial^2 f}{\partial y^2} \right]_0 \quad (16a)$$

$$[\lambda_\pi]_{y_\pi} = -2C \left[k + \frac{\partial f}{\partial y} \right]_{y_\pi}; \quad \left[\frac{\partial \lambda_\pi}{\partial y} \right]_{y_\pi} = -2C \left[\frac{\partial^2 f}{\partial y^2} \right]_{y_\pi} \quad (16b)$$

At a stable point $y_\pi = y_s$, the effective 4D potential (9) has a vanishing first derivative

$$\left[\frac{\partial V_{eff}}{\partial y_\pi} \right]_{y_\pi=y_s} = -\frac{e^{-4A(y_\pi)}}{C} \left[k + \frac{\partial f}{\partial y} \right]_{y=y_s}^2 = 0 \quad (17)$$

which at once implies that the second derivative $\partial^2 V_{eff} / \partial y_\pi^2$ also vanishes at the stable point y_s . The third derivative, however, is given at $y_\pi = y_s$ as

$$\left[\frac{\partial^3 V_{eff}}{\partial y_\pi^3} \right]_{y_\pi=y_s} = -\frac{2e^{-4A(y_s)}}{C} \left[\frac{\partial^2 f}{\partial y^2} \right]_{y=y_s}^2 \quad (18)$$

which is, of course, non-vanishing in consistence with Eq.(15a).

Considering, for example

$$f = a_1 \exp(-2a_2 y) \quad (19)$$

we find from Eqs.(15)

$$\phi_{cl} = \sqrt{a_1} e^{-a_2 y}; \quad V = \frac{1}{8} \left[\frac{\partial W}{\partial \phi} \right]^2 - 2CW^2 \quad (20)$$

where $W = 12kM^3 - a_2\phi^2$. Clearly, one gets back the same results as discussed in [13].

We now consider a tachyon-like scalar in the bulk [17, 18]. The action is given as

$$S = - \int d^5x \sqrt{-G} \left[R + V \sqrt{1 - \partial_A \phi \partial^A \phi} \right] - \int d^4x \sqrt{-g_i} \lambda_i(\phi) \quad (21)$$

where V is the potential of the field.

With a similar metric ansatz the field equations take the form

$$A'^2 = -\frac{1}{12} \frac{V}{\sqrt{1 + \phi'^2}}; \quad A'' = \frac{1}{6} \frac{V\phi'^2}{\sqrt{1 + \phi'^2}} \quad (22a)$$

$$\frac{\phi''}{(1 + \phi'^2)} - 4A'\phi' - \frac{1}{V} \frac{\partial V}{\partial \phi} = 0 \quad (22b)$$

and the corresponding boundary conditions are

$$[A']_i = \frac{1}{6} \lambda_i(\phi); \quad \left[\frac{V\phi'}{(1 + \phi'^2)^{3/2}} \right]_i = \frac{\partial \lambda_i(\phi)}{\partial \phi} \quad (23)$$

where $\{i\}$ corresponds to the brane locations at orbifold fixed points along the extra direction.

Proceeding along the same line as in the previous section, we obtain from Eq.(22)

$$\phi'^2 = -\frac{A''}{2A'^2}; \quad V = -12A' \sqrt{A'^2 - A''/2}. \quad (24)$$

Let us now consider the simple solution ansatz as

$$A = ky + a_1(y - a_2)^2 \quad (25)$$

where k, a_1, a_2 are the parameters of the theory. We get

$$\phi_{cl}'^2 = -\frac{a_1}{[y + 2a_1(y - a_2)]^2} \quad (26a)$$

$$V = -12[k + 2a_1(y - a_2)] \sqrt{1 - \frac{a_1}{k + 2a_1(y - a_2)}}. \quad (26b)$$

These equations imply that $a_1 = 0$ gives $V = -12k$ and $\phi = \text{constant}$, indicating pure RS background. We are, however, interested in $a_1 \neq 0$, in which case a real solution for ϕ can be obtained only when a_1 is negative:

$$\phi_{cl} = \pm (-4a_1)^{-1/2} \ln[k + 2a_1(y - a_2)]. \quad (27)$$

After a long but straightforward calculation following GW [8], we get

$$\frac{\partial V_{eff}}{\partial y_\pi} = -72e^{-4A} \left[\frac{-A''^2 + 4A'^4 - A''A'^2}{12A'^2 - 6A''} \right]_{y=y_\pi}. \quad (28)$$

So the stable point is at

$$y_s = \sqrt{\frac{1}{2|a_1|(\sqrt{5} + 1)}} + \frac{k}{2|a_1|} + a_2. \quad (29)$$

The corresponding metric function, upon setting $|a_1| = m^2$, is given by

$$A = \frac{k^2}{4m^2} + ka_2 - 0.154 \quad (30)$$

It is now clear that one can easily get the acceptable hierarchy by appropriately choosing the values of k/m and ka_2 with the stabilized radius y_s .

Our work clearly reveals an inherent instability in Randall-Sundrum two brane model. It is shown that just by introducing a canonical scalar field in the bulk, as initially suggested by GW, stabilization can not be achieved even with an arbitrary potential if full back-reaction of the scalar field on the metric is considered. However, we also show that in such models the two-brane separation can indeed be stabilized if the bulk scalar field has a tachyon-like action with respect to the bulk coordinate. We finally show that the full back-reacted metric in such case can also yield the desired warping from Planck scale to TeV scale and thus resolves the fine tuning problem of the Higgs mass in a stable braneworld scenario.

Acknowledgment

DM acknowledges Council of Scientific and Industrial Research, Govt. of India for providing financial support.

-
- [1] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. **B429** 263 (1998); I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. **B436** 257 (1998).
 [2] I. Antoniadis, Phys. Lett. **B246** 377 (1990); J. D. Lykken, Phys. Rev. **D54** 3693 (1996); R. Sundrum, Phys.

- Rev. **D59** 085009 (1999); K. R. Dienes, E. Dudas and T. Gherghetta, Phys. Lett. **B436** 55 (1998); G. Shiu and S. H. Tye, Phys. Rev. **D58** 106007 (1998); Z. Kakushadze and S. H. Tye, Nucl. Phys. **B548** 180 (1999).
 [3] P. Horava and E. Witten, Nucl. Phys. **B475** 94 (1996); *ibid* Nucl. Phys. **B460** 506 (1996).

- [4] L. Randall and R. Sundrum, Phys. Rev. Lett. **83** 3370 (1999); *ibid* Phys. Rev. Lett. **83** 4690 (1999).
- [5] N. Arkani-Hamed, S. Dimopoulos, G. Dvali and N. Kaloper, Phys. Rev. Lett. **84** 586 (2000); J. Lykken and L. Randall, JHEP **06** 014 (2000); C. Csaki and Y. Shirman, Phys. Rev. **D61** 024008 (2000); A. E. Nelson, Phys. Rev. **D63** 087503 (2001).
- [6] A. G. Cohen and D. B. Kaplan, hep-th/9910132; C. P. Burgess, L. E. Ibanez and F. Quevedo, Phys. Lett. **B447** 257 (1999); A. Chodos and E. Poppitz, Phys. Lett. **B471** 119 (1999); T. Gherghetta and M. Shaposhnikov, Phys. Rev. Lett. **85** 240 (2000).
- [7] N. Kaloper, Phys. Rev. **D60** 123506 (1999); T. Nihei, Phys. Lett. **B465** 81 (1999); H. B. Kim and H. D. Kim, Phys. Rev. **D61** 064003 (2000).
- [8] W. D. Goldberger and M. B. Wise, Phys. Rev. Lett. **83** 4922 (1999); *ibid* Phys. Lett. **B475** 275 (2000).
- [9] M. Luty and R. Sundrum, Phys. Rev. **D64** 065012 (2001); F. Brummer, A. Hebecker and E. Trincherini, Nucl. Phys. **B738** 283 (2006); G.L. Alberghi *et al*, Phys. Rev. **D72** 025005 (2005); T. Kobayashi and K. Yoshioka, JHEP **0411** 024 (2004); S. Ichinose and A. Murayama, Phys. Lett. **B625** 106 (2005); S. Kalyana Rama, Phys. Lett. **B495** 176 (2000); H. K. Jassal, hep-th/0312253.
- [10] N. Maru, N. Okada, hep-th/0508113; M. Eto, N. Maru and N. Sakai, Phys. Rev. **D70** 086002 (2004); N. Maru and N. Okada, Phys. Rev. **D70** 025002 (2004);
- [11] S. Das, A. Dey, S. SenGupta, hep-th/0511247; H. Yoshiguchi *et al*, JCAP **0603** 018 (2006); E. E. Boos *et al*, hep-th/0511185; R. Maartens, Living Rev. Rel. **7** 7 (2004) and references therein.
- [12] I. I. Kogan *et al*, Nucl. Phys. **B584** 313 (2000); D. Choudhury *et al*, JHEP **09** 021 (2000); S. Forste, hep-th/0110055.
- [13] O. DeWolfe, D. Z. Freedman, S. S. Gubser and A. Karch, Phys. Rev. **D62** 046008 (2000).
- [14] C. Csaki, M. L. Graesser and G. D. Kribs, Phys. Rev. **D63** 065002 (2001); C. Csaki, M. L. Graesser, L. Randall and J. Terning, Phys. Rev. **D62** 045015 (2000).
- [15] W. D. Goldberger and M. B. Wise, Phys. Rev. **D60** 107505 (1999); S. Kachru, M. B. Schulz and E. Silverstein, Phys. Rev. **D62** 045021 (2000); H. A. Chamblin and H. S. Reall, Nucl. Phys. **B562** 133 (1999); C. Csaki, hep-ph/0404096; R. Neves, TSPU Vestnik **44N7** 94 (2004); E. Dudas and M. Quiros, Nucl. Phys. **B721** 309 (2005).
- [16] K. Behrndt and M. Cvetič, Phys. Lett. **B475** 253 (2000).
- [17] A. Sen, JHEP **0204**, 048 (2002); *ibid* JHEP **0207**, 065 (2002); *ibid* Mod. Phys. Letts. **A17**, 1797 (2002).
- [18] R. Koley and S. Kar, Phys. Lett. **B623** 244 (2005); M. Pospelov, hep-ph/0412280.